

Spontaneous Heterotopic Pregnancy: A Case Report and Review of Radiological Assessment and Diagnostic Challenges

Biswas SS*

Consultant Radiologist, Pragathi Teleradiology, India

ABSTRACT

Heterotopic pregnancy is the simultaneous occurrence of intrauterine and extrauterine gestation. It is a rare complication of gestation more common when conception is achieved by assisted reproduction techniques than in natural conceptions. Heterotopic pregnancy is a life-threatening pathology with the diagnosis often missed radiological investigations due to its rarity. A painstaking sonographic evaluation is required to rule out the presence of heterotopic gestation and thus help in prompt diagnosis and appropriate management. Early diagnosis of this pathology reduces the incidence of complications as well as maternal mortality. We hereby report a case of heterotopic pregnancy presenting with gestation amenorrhea of seven weeks and acute pain in the lower abdomen associated with episodes of spotting per vagina of three days duration. The diagnosis of heterotopic pregnancy with ruptured extrauterine gestation was made on a careful ultrasound examination and was managed with exploratory laparotomy. The intrauterine pregnancy course was uneventful with a delivery of a live baby at term. This case report emphasizes the need to carefully examine the adnexa even with the visualization of normal pregnancy on ultrasound interrogation to rule out heterotopic pregnancy.

Keywords: Heterotopic pregnancy, Extrauterine pregnancy, Amenorrhoea, Exploratory laparotomy.

INTRODUCTION

Heterotopic pregnancy is the coexistence of intrauterine and extrauterine gestations [1,2].

It is a rare and life-threatening situation that is difficult to diagnose and easily missed [3]. The estimated incidence of heterotopic pregnancy was 1 in 30,000 in natural conceptions. However, in recent years due to the increasing incidence of pelvic inflammatory diseases and rising use of intrauterine contraceptive devices, the incidence of heterotopic has markedly increased to around 1:7,000 in the general population [2,4]. The incidence of heterotopic pregnancy is even higher with assisted conceptions [4].

Vol No: 07, Issue: 03

Received Date: November 29, 2022

Published Date: December 07, 2022

*Corresponding Author

Biswas SS

Consultant Radiologist, Pragathi Teleradiology,
Hyderabad, Telangana, India

E-mail: ssbinmemphis@gmail.com

Citation: Biswas SS. (2022). Spontaneous Heterotopic Pregnancy: A Case Report and Review of Radiological Assessment and Diagnostic Challenges. Mathews J Case Rep. 7(3):76.

Copyright: Biswas SS © (2022). This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

The most frequent ectopic site is the fallopian tube in both natural and assisted conceptions while the cornua are the second most common location. The other less common sites of ectopic gestations are the ovary, the cervix, and the abdomen [4,5]. More than half of the intrauterine gestations in heterotopic gestations result in live deliveries if detected early and managed appropriately [2,4]. A delay or missed diagnosis can lead to a rise in morbidity and mortality both for the mother and the intrauterine gestation.

Herein, we report a case of ruptured heterotopic pregnancy presenting at 7 weeks of gestation with lower abdominal pain and bleeding per vagina which was managed with immediate exploratory laparotomy. The intrauterine pregnancy course was uneventful with spontaneous vaginal delivery of a healthy baby at term.

CASE REPORT

She was a 39-year-old multigravida ($G_4P_2^{+1}$, 2 alive), who presented to our hospital with a history of cessation of menses for seven weeks, progressive lower abdominal pain, and bleeding per vagina of three days duration. There was no preceding history of pelvic inflammatory disease or fertility treatment. No family history of twins was present. Personal history had no significant findings. Her blood pressure on admission was 90/55mmHg, her pulse rate was 105/minute, and a respiratory rate of 20/minute. There was suprapubic tenderness and mild distension on abdomen examination. She had an enlarged uterus corresponding to seven weeks size of gestation. The cervix was closed with a tender left adnexa. Her hemoglobin level was 8.0 gm/dl with a positive beta-hCG test. The transabdominal ultrasonography showed a moderate amount of fluid in the peritoneal cavity and the pelvic region with a viable intrauterine gestation of about 7 weeks and a left adnexal haematoma of size 11 X 7 cm with a central, dense echogenic ring indicative suggestive of ruptured left ectopic gestation (Figure 1).




Figure 1: A live embryo within an intrauterine Gestational Sac (GS) with a left adnexal mass and surrounding hematoma on longitudinal view of an abdominopelvic ultrasound scan.

A diagnosis of heterotopic pregnancy with ruptured left-sided ectopic gestation was made in view of clinical history and sonographic findings. The patient underwent an emergency exploratory laparotomy. The intra-operative findings were a ruptured left-sided tubal pregnancy and hemoperitoneum

of 1.4 litres. A left-sided salpingectomy was performed and the patient was transfused with three units of blood intra-operatively. The intrauterine live gestation was allowed to progress (Figure 2) with the patient delivering a healthy live baby at 39 weeks.




Figure 2: Normal left adnexa with a viable intrauterine gestation 10 weeks after surgery.

DISCUSSION

Heterotopic gestation is defined as the co-existence of both intrauterine and extrauterine pregnancy. It frequently occurs in the fallopian tube and rarely in the cervix and ovary [1,4]. Though fairly common with assisted reproduction, heterotopic gestation rarely occurs in natural conceptions [2,4]. Triplet and quadruplet heterotopic conceptions have also been reported, though the incidences are extremely rare [4], Heterotopic pregnancy is seen in 1 in 30,000 natural conceptions, 1 in 900 drug-induced pregnancies, and higher in assisted conceptions [4,5].

The risk factors for heterotopic pregnancy include previous tubal surgeries, previous ectopic pregnancies, and assisted reproduction techniques like in vitro fertilization and pharmacological ovulation induction [4,5]. This index patient had no associated risk factors for heterotopic conception.

The diagnosis of heterotopic pregnancy is difficult to make clinically [2,4,5] According to Tal, et al. [5] the principal presenting symptoms, in decreasing frequency, are amenorrhoea, abdominal pain, vaginal bleeding, abdominal swelling, and shock. In this case report, the patient presented with these symptoms and was in shock from the hemoperitoneum due to the ruptured tubal pregnancy. Heterotopic pregnancy may likely be missed in spontaneous conceptions unless the sonologist meticulously examines the fallopian tubes and the pelvis. If overlooked, it may lead to tubal rupture and acute abdominal syndrome as in this index case and can progress to maternal shock leading to maternal mortality [4,5].

Early diagnosis of heterotopic pregnancy is challenging as the detection of an intrauterine implanted embryo and positive beta-hCG test can mask the need to carefully scan the adnexa in asymptomatic patients [5]. About half of heterotopic pregnancies are discovered during emergency exploratory laparotomies for ruptured ectopic gestation [2].

In this index case, intrauterine and extrauterine pregnancies were discovered via an emergency ultrasound scan. A high-resolution ultrasonography with focused adnexal scanning at four to six weeks of gestation is recommended for early detection of heterotopic pregnancy [4,5].

Certain pathology often mimic heterotopic gestation and can result in delayed or missed diagnosis. Intrauterine gestation with hemorrhagic corpus luteum can simulate heterotopic pregnancy both clinically and on ultrasound [6]. Bicornuate uterus with pregnancy in both horns can also mimic a heterotopic pregnancy. A high-resolution ultrasound scan with color Doppler is helpful in ruling out these differential diagnoses as heterotopic gestation displays increased blood flow with markedly reduced resistance index. This is an important aid in the diagnosis of heterotopic pregnancy.

The management of heterotopic pregnancy is laparoscopy or laparotomy for tubal pregnancy [8]. However, factors such as the severity of presentation, site of ectopic gestation, number of previous pregnancies, the viability or otherwise of the intrauterine pregnancy, the expertise of the physicians, and the socioeconomic status of the patients should be taken into consideration in deciding the options of management [2,4,5]. Usually, the least invasive procedures are preferred for a better outcome for the intrauterine pregnancy. Laparoscopic surgery (salpingostomy or salpingectomy) is the treatment of choice in patients with haemodynamic stability because of a better outcome for intrauterine pregnancy and the least harmful effects on intrauterine gestation. Laparotomy should be carried out in patients with serious intra-abdominal bleeding or in patients with haemodynamic instability [9]. Our patient presented with a rupture of ectopic pregnancy with hemoperitoneum. Exploratory laparotomy was carried out to remove the extrauterine non-viable gestation while allowing the viable intrauterine pregnancy to develop to term. Thus, all surgeons operating on ruptured ectopic must bear in mind the likelihood of heterotopic pregnancy

and must handle the uterus with care [10]. Hegazy (2014) stated that pregnancy in twins has a familial tendency [11]. However, in our case, there was no family history and personal history was also not significant.

CONCLUSION

Early diagnosis of heterotopic pregnancy significantly reduces the morbidity and mortality of both the mother and the intrauterine gestation. This case report emphasizes focused scanning of the adnexa in the first trimester to exclude extrauterine gestation and to rule out heterotopic pregnancy.

CONSENT AND ETHICAL APPROVAL

We declared that we have obtained all required consent. The patient has also agreed for her clinical images and information to be published in the medical journal. She understands that her identity will not be revealed.

REFERENCES

1. Govindarajan MJ, Rajan R. (2008). Heterotopic pregnancy in natural conception. *J Hum Reprod Sci.* 1:37-38.
2. Nabi U, Yousaf A, Ghaffar F, Sajid S, Ahmed MMH. (2019). Heterotopic Pregnancy - A Diagnostic Challenge. Six Case Reports and Literature Review. *Cureus.* 11(11):e6080.
3. Hassani KI, Bouazzaoui AE, Khatouf M, Mazaz K. (2010). Heterotopic pregnancy: a diagnosis we should suspect more often. *J Emerg Trauma Shock.* 3(3):304.
4. Reece EA, Petrie RH, Sirmans MF, Finster M, Todd WD. (1983). Combined intrauterine and extrauterine gestations: a review. *Am J Obstet Gynecol.* 146(3):323-30
5. Tal J, Haddad S, Gordon N, Timor-Tritsch I. (1996). Heterotopic pregnancy after ovulation induction and assisted reproductive technologies: a literature review from 1971 to 1993. *Fertil Steril.* 66(1):1-12.
6. Nguyen-Tran C, Toy EC. (2000). Heterotopic pregnancy: viable twin intrauterine pregnancy with a viable right tubal ectopic pregnancy. *J Ultrasound Med.* 19(5):355.
7. Jaffe R, Warsof SL. (1991). Transvaginal color Doppler imaging in the assessment of uteroplacental blood flow in the normal first-trimester pregnancy. *Am J Obstet Gynecol.* 164(3):781-785.
8. Gruber I, Lahodny J, Illmensee K, Losch A. (2002). Heterotopic pregnancy: Report of three cases. *Wien Klin Wochenschr.* 114:229-232.
9. Melendez J, Paraskevopolou SM, Foo X, Yoong W. (2010). Heterotopic pregnancy: tubal ectopic pregnancy with a viable IVF intrauterine pregnancy. *J Obstet Gynaecol.* 30:742-743.
10. Sohail S. (2005). Haemorrhagic corpus luteum mimicking heterotopic pregnancy. *J Coll Physicians Sur Pak.* 15:180-181.
11. Hegazy A. (2014). *Clinical embryology for medical students and postgraduate doctors.* Lap Lambert Academic Publishing:184.